

WEEDS	WHEAT	BARLEY	OATS
Annual Smartweed	●	●	●
Ball Mustard	●	●	●
Chickweed (1)	●	●	●
Corn Spurry	●	●	●
Cow Cockle	●	●	●
Flixweed	●	●	●
Green Smartweed	●	●	●
Groundsel	●	●	●
Hemp-Nettle	●	●	●
Kochia*	●	●	●
Lady's Thumb	●	●	●
Lamb's Quarters	●	●	●
Narrow-Leaved Hawk's Beard	●	●	●
Redroot Pigweed	●	●	●
Russian Thistle	●	●	●
Shepherd's Purse	●	●	●
Stinkweed	●	●	●

WEEDS	WHEAT	BARLEY	OATS
Tartary Buckwheat	●	●	●
Volunteer Rapeseed (2)	●	●	●
Volunteer Sunflower	●	●	●
Wild Buckwheat (3)	●	●	●
Wild Mustard	●	●	●
Canada Thistle (4)	/	/	/
Cleavers	/	/	/
Round Leaved Mallow (5)	/	/	/
Scentless Chamomile	/	/	/
Sow Thistle (4)	/	/	/
Stork's Bill	/	/	/
Toadflax (6)	/	/	/

● Control / Suppression

- (1) Apply when chickweed is small (1-6 leaf) and actively growing, but before crop canopy prevents thorough herbicide coverage of weeds. Chickweed emerging after application will not be controlled.
- (2) MPOWER R alone will not control imazamox and imazethapyr tolerant canola (e.g., Clearfield canola).
- (3) Apply to actively growing wild buckwheat in the cotyledon 1-3 leaf stage. Under environmental stress, such as dry weather, control may be reduced. Large plants may re-grow after treatment.
- (4) Apply when the majority of thistles have emerged and are actively growing
- (5) Apply to actively growing round leaved mallow from the 2-6 leaf stage (10-12 cm in height).
- (6) Apply to toadflax that is no larger than 15 cm in height.
- \*Not including Group 2 resistant bio-type