

PMRA Approved Label – M.A.
2021-4039 2021-09-27

GROUP	1	HERBICIDE
-------	---	-----------

AURORA® HERBICIDE

COMMERCIAL (AGRICULTURAL)

EMULSIFIABLE CONCENTRATE

A post emergence herbicide for control of wild oats, volunteer (tame) oats, green foxtail, yellow foxtail, barnyard grass, Persian dandelion and volunteer canary seed in Spring wheat and Durum wheat.

FOR SALE FOR USE IN THE PRAIRIE PROVINCES AND INTERIOR OF BRITISH COLUMBIA (INCLUDING THE PEACE RIVER REGION), OKANAGAN AND CRESTON FLATS REGIONS OF BRITISH COLUMBIA ONLY.

ACTIVE INGREDIENT: Clodinafop-propargyl 240 g/L
Warning: contains the allergen epoxidized soybean oil

REGISTRATION NO.: 29711 PEST CONTROL PRODUCTS ACT

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING.

KEEP OUT OF REACH OF CHILDREN.

CAUTION



POISON

WARNING: EYE & SKIN IRRITANT

NET CONTENTS: 1.84L – bulk

For Hazardous Materials [or Dangerous Goods] Incident, Spill, Leak, Fire, Exposure, or Accident Call CHEMTREC 1-800-424-9300/ +1-703-527-3887

NewAgco, Inc.
320 22nd Street East
Saskatoon (Sask.)
S7K 0H1 Canada
1-844-269-3276

Distributed by:
AgraCity Crop & Nutrition Ltd
320-22nd Street East
Saskatoon (Sask.)
S7K 0H1 Canada
1-844-269-3276

AURORA®, CANUCK®, CLOBBER®, FOXXY™, and MPOWER® are trademarks of AgraCity Crop & Nutrition Ltd. or its affiliates.

PMRA Approved Label – M.A.

2021-4039 2021-09-27

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

FIRST AID

IN CASE OF POISONING, contact a physician or a poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If on skin or clothing, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If swallowed, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

TOXICOLOGICAL INFORMATION

This product contains a PETROLEUM DISTILLATE. DO NOT INDUCE VOMITING. Vomiting may cause aspiration pneumonia. Treat symptomatically for ingestion and/or skin and eye contact.

PRECAUTIONS

Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

- **CAUTION – POISON**
- **WARNING – EYE AND SKIN IRRITANT**
- **KEEP OUT OF THE REACH OF CHILDREN.**
- Harmful if swallowed.
- May irritate eyes. Do not wear contact lenses when using.
- DO NOT get in eyes or on skin. Avoid contact with clothing.
- During mixing, loading, clean-up, repair and when applying by groundboom, workers must wear coveralls over long-sleeved shirt and long pants, shoes and socks, goggles, and chemical-resistant gloves. When applying by aerial application, pilots must wear long-sleeved shirt and long pants.
- Wash gloves thoroughly with soap and water before removing them during any operation.
- Wash hands thoroughly with soap and water after using this product and before eating, drinking or smoking.
- Remove contaminated clothing immediately after use. Store and wash contaminated clothing separately from household laundry before reuse. Wash thoroughly with soap and water after handling. Handle and apply only as recommended on this label.
- Do not eat, drink or smoke while mixing, loading or during application.
- DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PMRA Approved Label – M.A.
2021-4039 2021-09-27

- If mixing and loading more than 15 kg a.i. in a day, workers must use a closed mixing/loading system

ENVIRONMENTAL PRECAUTIONS

This product contains an active ingredient and aromatic petroleum distillates which are toxic to aquatic organisms.

TOXIC to aquatic organisms and non-target terrestrial plants. Observe buffer zones specified under DIRECTIONS FOR USE.

ENVIRONMENTAL PRECAUTIONS

- TOXIC to non-target terrestrial plants. Observe buffer zones specified under DIRECTIONS FOR USE.
- TOXIC to aquatic organisms. Observe buffer zones specified under DIRECTIONS FOR USE.
- The residues of this product demonstrate the properties and characteristics associated with chemicals detected in ground water. The use of clodinafop-propargyl products in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.
- To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil or clay.
- Avoid application when heavy rain is forecast.
- Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

When a tank mixture is used, consult the labels of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in U.S., visit CropLife Canada's website at www.croplife.ca.

STORAGE

To prevent contamination, store this product away from food or feed.

DISPOSAL OF UNUSED, UNWANTED PRODUCT

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill or for clean-up of spills.

CONTAINER DISPOSAL:

For recyclable containers:

Do not reuse this container for any purpose. This is a recyclable container, and may be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

- 1 Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
- 2 Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial

PMRA Approved Label – M.A.
2021-4039 2021-09-27
requirements.

For refillable containers: For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not use this container for any other purpose.

PMRA Approved Label – M.A.

2021-4039 2021-09-27

{LABEL BOOKLET}

GROUP

1

HERBICIDE

AURORA® HERBICIDE

COMMERCIAL (AGRICULTURAL)

EMULSIFIABLE CONCENTRATE

A post emergence herbicide for control of wild oats, volunteer (tame) oats, green foxtail, yellow foxtail, barnyard grass, Persian darnel and volunteer canary seed in Spring wheat and Durum wheat.

FOR SALE FOR USE IN THE PRAIRIE PROVINCES AND INTERIOR OF BRITISH COLUMBIA (INCLUDING THE PEACE RIVER REGION), OKANAGAN AND CRESTON FLATS REGIONS OF BRITISH COLUMBIA ONLY.

ACTIVE INGREDIENT: Clodinafop-propargyl 240 g/L

Warning: contains the allergen epoxidized soybean oil

REGISTRATION NO.: 29711 PEST CONTROL PRODUCTS ACT

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING.

KEEP OUT OF REACH OF CHILDREN.

CAUTION



POISON

WARNING: EYE & SKIN IRRITANT

NET CONTENTS: 1.84 L - Bulk

For Hazardous Materials [or Dangerous Goods] Incident, Spill, Leak, Fire, Exposure, or Accident Call CHEMTREC 1-800-424-9300/ +1-703-527-3887

NewAgco, Inc.
320 22nd Street East
Saskatoon (Sask.)
S7K 0H1 Canada
1-844-269-3276

Distributed by:
AgraCity Crop & Nutrition Ltd
320-22nd Street East
Saskatoon (Sask.)
S7K 0H1 Canada
1-844-269-3276

AURORA®, CANUCK®, CLOBBER®, FOXXY™, and MPOWER® are trademarks of AgraCity Crop & Nutrition Ltd. or its affiliates.

PMRA Approved Label – M.A.

2021-4039 2021-09-27

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

FIRST AID

IN CASE OF POISONING, contact a physician or a poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If on skin or clothing, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If swallowed, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

TOXICOLOGICAL INFORMATION

This product contains a PETROLEUM DISTILLATE. DO NOT INDUCE VOMITING. Vomiting may cause aspiration pneumonia. Treat symptomatically for ingestion and/or skin and eye contact.

PRECAUTIONS

Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

- CAUTION – POISON
- WARNING – EYE AND SKIN IRRITANT
- KEEP OUT OF THE REACH OF CHILDREN.
- Harmful if swallowed.
- May irritate eyes. Do not wear contact lenses when using.
- DO NOT get in eyes or on skin. Avoid contact with clothing.
- During mixing, loading, clean-up, repair and when applying by groundboom, workers must wear coveralls over long-sleeved shirt and long pants, shoes and socks, goggles, and chemical-resistant gloves. When applying by aerial application, pilots must wear long-sleeved shirt and long pants.
- Wash gloves thoroughly with soap and water before removing them during any operation.
- Wash hands thoroughly with soap and water after using this product and before eating, drinking or smoking.
- Remove contaminated clothing immediately after use. Store and wash contaminated clothing separately from household laundry before reuse. Wash thoroughly with soap and water after handling. Handle and apply only as recommended on this label.
- Do not eat, drink or smoke while mixing, loading or during application.
- DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of

PMRA Approved Label – M.A.

2021-4039 2021-09-27

12 hours.

- If mixing and loading more than 15 kg a.i. in a day, workers must use a closed mixing/loading system.

ENVIRONMENTAL PRECAUTIONS

This product contains an active ingredient and aromatic petroleum distillates which are toxic to aquatic organisms.

TOXIC to aquatic organisms and non-target terrestrial plants. Observe buffer zones specified under DIRECTIONS FOR USE.

ENVIRONMENTAL PRECAUTIONS

- **TOXIC** to non-target terrestrial plants. Observe buffer zones specified under DIRECTIONS FOR USE.
- **TOXIC** to aquatic organisms. Observe buffer zones specified under DIRECTIONS FOR USE.
- The residues of this product demonstrate the properties and characteristics associated with chemicals detected in ground water. The use of clodinafop-propargyl products in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.
- To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil or clay.
- Avoid application when heavy rain is forecast.
- Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

When a tank mixture is used, consult the labels of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in U.S., visit CropLife Canada's website at www.croplife.ca.

STORAGE

To prevent contamination, store this product away from food or feed.

DISPOSAL OF UNUSED, UNWANTED PRODUCT

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill or for clean-up of spills.

PMRA Approved Label – M.A.
2021-4039 2021-09-27

CONTAINER DISPOSAL:

For recyclable containers:

Do not reuse this container for any purpose. This is a recyclable container and may be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For refillable containers: For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not use this container for any other purpose.

PRODUCT INFORMATION

AURORA HERBICIDE is a systemic, post-emergence herbicide for the selective control of wild oats, volunteer (tame) oats, green foxtail, yellow foxtail, barnyard grass, Persian dandelion and volunteer canary seed in all types of Spring wheat and Durum wheat. Do not use **AURORA HERBICIDE** on barley, as crop injury will occur. Do not apply this product using aerial application equipment except under conditions specified on this label.

AURORA HERBICIDE is absorbed by the leaves and is rapidly translocated to the growing points of leaves and stems. Thorough coverage of the plants is essential for consistent control. Actively growing susceptible grasses stop growing within 48 hours of treatment. Depending on species, growing conditions and crop competition, leaves and growing points turn yellow within one to three weeks after application. Further colour changes and loss of vigour will be observed, followed by a browning and complete control three to five weeks after application.

Although **AURORA HERBICIDE** does not control broadleaf weeds, **AURORA HERBICIDE** can be tank-mixed with a wide range of broadleaf herbicides to provide broad spectrum weed control in wheat. See section entitled "TANK MIXES of **AURORA HERBICIDE** WITH BROADLEAF WEED HERBICIDES" and refer to the appropriate Tank Mix section for ground and aerial application.

DIRECTIONS FOR USE

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

A 30-day plant-back interval must be observed for all unlabelled crops.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Field sprayer application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American

PMRA Approved Label – M.A.
2021-4039 2021-09-27

Society of Agricultural Engineers (ASAE S572.1) medium classification. Boom height must be 60 cm or less above the crop or ground.

Aerial application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply when wind speed is greater than 16 km/h at flying height at the site of application. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) coarse classification. Reduce drift caused by turbulent wingtip vortices. The nozzle distribution along the spray boom length **MUST NOT** exceed 65% of the wing- or rotorspan.

CROPS: Spring Wheat and Durum Wheat

AURORA HERBICIDE can be used on all varieties of spring wheat and Durum wheat. Observe minimum interval to harvest of 60 days after treatment.

Do not apply on barley or any crop other than Spring Wheat or Durum wheat, as crop damage will result. Do not allow spray to drift to adjacent fields seeded to crops other than spring wheat or Durum wheat.

Do not treat wheat underseeded to forages.

Observe a minimum pre-harvest interval of 60 days after treatment for grain and straw and of 30 days after treatment for hay. Observe a minimum of seven (7) days before grazing livestock on treated crops.

For use directions specific to application by air, please refer to sections on aerial application. For aerial application precautions, please refer to the precautions section.

WEEDS CONTROLLED: Wild Oats, Volunteer (Tame) Oats, Green Foxtail, Yellow Foxtail, Barnyard grass, Persian Darnel and Volunteer Canary Seed

TIMING OF APPLICATION:

TIMING	GROWTH STAGE	ADDITIONAL REMARKS
WILD OATS	1 to 6 leaf stage on main stem	Prior to emergence of 4th tiller
VOLUNTEER (TAME) OATS	3 to 6 leaf stage on main stem	Prior to emergence of 4th tiller
GREEN FOXTAIL AND YELLOW FOXTAIL (wild millet, pigeon grass)	1 to 5 leaf stage on main stem	For optimum control apply prior to emergence of the 3 rd tiller and while foxtail is actively growing
BARNYARD GRASS	1 to 5 leaf stage on main stem	For optimum control apply before tillering, and while barnyard grass is actively growing
PERSIAN DARNEL	1 to 5 leaf stage on main stem	For optimum control apply before tillering, and while Persian darnel is actively growing
VOLUNTEER CANARY SEED	1 to 6 leaf stage on main stem	Prior to emergence of 4th tiller
SPRING WHEAT AND DURUM WHEAT	Prior to emergence of 4th tiller	When tank-mixing with a broadleaf herbicide, always refer to the label of the broadleaf partner prior to use.

- For optimum results, apply **AURORA HERBICIDE** to actively growing weeds. An early application will maximize crop yields by reducing weed competition. Weeds emerging after application of **AURORA HERBICIDE** will not be controlled.

PMRA Approved Label – M.A.
2021-4039 2021-09-27

- Weed control following application of **AURORA HERBICIDE** alone, or in combination with broadleaf weed herbicides, can be reduced or delayed under stress conditions such as drought, heat, insufficient fertility, flooding or prolonged cool temperatures. Grass escapes or re-tillering may occur if application is made during prolonged stress conditions. Optimum weed control will be obtained if application of **AURORA HERBICIDE** is delayed until the stress conditions have ended and weeds are once again actively growing.
- **AURORA HERBICIDE** alone can be used 30 minutes before rainfall.
- Do not apply to crop that is stressed by conditions such as frost, low fertility, drought, flooding, disease or insect damage as crop injury may result.

RATE OF USE FOR GROUND APPLICATION:

Apply the recommended rate of **AURORA HERBICIDE** and the recommended rate of an adjuvant of either SCORE®, ASSIST®, or CHEM SPRAY® Crop Oil Concentrate in a minimum of 50 to 100 L of water per hectare.

<p>To control:</p> <p>WILD OATS VOLUNTEER (TAME) OATS GREEN FOXTAIL YELLOW FOXTAIL BARNYARD GRASS VOLUNTEER CANARY SEED</p>	<p>To control:</p> <p>WILD OATS VOLUNTEER (TAME) OATS GREEN FOXTAIL YELLOW FOXTAIL BARNYARD GRASS VOLUNTEER CANARY SEED PERSIAN DARNEL</p>
<p>Rates for use with 50 L/ha water</p>	
<p>Apply:</p> <p>230 mL/ha of AURORA HERBICIDE + 400 mL/ha of (SCORE, ASSIST, or CHEM SPRAY Crop Oil Concentrate) with 50 L/ha of water (0.8% volume/volume)</p>	<p>Apply:</p> <p>290 mL/ha of AURORA HERBICIDE + 500 mL/ha of (SCORE or ASSIST, or CHEM SPRAY Crop Oil Concentrate) with 50 L/ha of water (1.0% volume/volume)</p>
<p>NOTE: One 1.84/3.68/4.0/4.7/8.0 L jug of AURORA HERBICIDE + 3.2/6.4/6.96/8.16/14.0 L of (SCORE, ASSIST or CHEM SPRAY Crop Oil Concentrate) treats 8 ha (20 acres)/16 ha (40 acres)/17.4 ha (43 acres)/20.4 ha (50.4 acres)/35 ha (86.4 acres) with 400/800/870/1020/1750 3.68 L of water.</p>	<p>NOTE: One 1.84/3.68/4.0/4.7/8.0 L jug of AURORA HERBICIDE + 3.2/6.4/6.96/8.16/14.0 L of (SCORE, ASSIST or CHEM SPRAY Crop Oil Concentrate) treats 6.4 ha (15.8 acres)/12.8 ha (31.4 acres)/13.9 ha (34.3 acres)/16.3 ha (40.2 acres)/28 ha (69 acres) with 320/640/695/815/1400 L of water.</p>
<p>Rates for use with 100 L/ha water</p>	
<p>Apply:</p> <p>230 mL/ha of AURORA HERBICIDE + 800 mL/ha of (SCORE, ASSIST or CHEM SPRAY Crop Oil Concentrate) with 100 L/ha of water (0.8% volume/volume)</p>	<p>Apply:</p> <p>290 mL/ha of AURORA HERBICIDE + 1 L/ha of (SCORE, ASSIST or CHEM SPRAY Crop Oil Concentrate with 100 L/ha of water (1.0% volume/volume)</p>
<p>NOTE: One 1.84/3.68/4.0/4.7/8.0 L jug of AURORA HERBICIDE + 6.4/12.8/13.9/16.3/28.0 L of (SCORE, ASSIST or CHEM SPRAY Crop Oil Concentrate) treats 8 ha (20 acres)/16 ha (40 acres)/17.4 ha (43 acres)/20.4 ha (50.4 acres)/35 ha (86.4 acres) with 800/1600/1740/2040/3500 L of water.</p>	<p>NOTE: One 1.84/3.68/4.0/4.7/8.0 L jug of AURORA HERBICIDE + 6.4/12.8/13.9/16.3/28.0 L of (SCORE, ASSIST or CHEM SPRAY Crop Oil Concentrate) treats 6.4 ha (15.8 acres)/12.8 ha (31.4 acres)/13.9 ha (34.3 acres)/16.3 ha (40.2 acres)/28 ha (69 acres) with 640/1280/1390/1630/2800 L of water.</p>

PMRA Approved Label – M.A.
2021-4039 2021-09-27

NOTE: Always use an adjuvant of either SCORE, ASSIST or CHEM SPRAY Crop Oil Concentrate with AURORA HERBICIDE.

RATE OF USE FOR AERIAL APPLICATION:

Apply the recommended rate of **AURORA HERBICIDE** and the recommended rate of an adjuvant of either SCORE, ASSIST, or CHEM SPRAY Crop Oil Concentrate in a minimum of 30 L of water per hectare according to the following table:

To Control: WILD OATS	To Control: WILD OATS, GREEN FOXTAIL, YELLOW FOXTAIL, PERSIAN DARNEL
Apply: 230 mL/ha of AURORA HERBICIDE + 240 mL/ha of SCORE, ASSIST, or CHEM SPRAY Crop Oil Concentrate with 30 L/ha of water (0.8% volume/volume)	Apply: 290 mL/ha of AURORA HERBICIDE + 300 mL/ha of SCORE, ASSIST, or CHEM SPRAY Crop Oil Concentrate with 30 L/ha of water (1.0% volume/volume)
NOTE: One 1.84/3.68/4.0/4.7/8.0 L jug of AURORA HERBICIDE + 1.92/3.84/4.18/4.9/8.4 L SCORE, ASSIST or CHEM SPRAY Crop Oil Concentrate treats 8 ha (20 acres)/16 ha (40 acres)/17.4 ha (43 acres)/20.4 ha (50.4 acres)/35 ha (86.4 acres) with 240/480/522/612/1050 L of water.	NOTE: One 1.84/3.68/4.0/4.7/8.0 L jug of AURORA HERBICIDE + 1.92/3.84/4.17/4.9/8.4 L SCORE, ASSIST or CHEM SPRAY Crop Oil Concentrate treats 6.4 ha (15.8 acres)/12.8 ha (31.4 acres)/13.9 ha (34.3 acres)/16.3 ha (40.2 acres)/28 ha (69 acres) with 192/384/417/490/840 L of water.

NOTE: Always use adjuvants of either SCORE, ASSIST or CHEM SPRAY Crop Oil Concentrate with AURORA HERBICIDE.

TANK MIXES of AURORA HERBICIDE WITH BROADLEAF WEED HERBICIDES – GROUND APPLICATION:

CROP: SPRING and DURUM WHEAT

TANK MIXES WITH BROADLEAF WEED HERBICIDES, IN SPRING WHEAT and DURUM WHEAT:

For broad spectrum control of wild oats, green foxtail and broadleaf weeds, **AURORA HERBICIDE** can be tank-mixed with broadleaf herbicides as described in the following tables. Consult the label of the tank-mix partner for a list of broadleaf weeds controlled, rates, timing, recropping restrictions, grazing interval restrictions, recommendations for specific weeds, directions for use and precautions and follow the more restrictive label. When tank-mixing always add the broadleaf herbicide(s) to the spray tank first; followed by **AURORA HERBICIDE**, with SCORE or ASSIST Adjuvant, or CHEM SPRAY Crop Oil Concentrate added last. For the appropriate rate of **AURORA HERBICIDE** with SCORE, ASSIST or CHEM SPRAY Crop Oil Concentrate, refer to the 'Rate of Use' section of the label.

Tank-Mix Partner	Product Rates	Crop Stage ¹
Dyvel [®]	1.25 L/ha	2 to 5 leaf
Refine [®] Extra ² or MPOWER [®] RX ² HERBICIDE	20 g/ha	2 leaf to flag leaf
Buctril [®] M or CANUCK [®] HERBICIDE	1.0 L/ha	2 leaf to flag leaf
Estaprop [®]	1.75 L/ha	4 leaf to early flag leaf (shot blade)
Turboprop [®] 600	1.75 L/ha	4 leaf to early flag leaf (shot

		blade)
Dicloroprop® D	1.75 L/ha	4 leaf to early flag leaf (shot blade)
Lontrel™ 360 or CLOBBER® HERBICIDE tank mixed with MCPA Ester (assume 500 series)	280 to 420 mL/ha tank mixed with 1.1 L/ha	3 leaf to flag leaf
Curtail™ M or CLOBBER M HERBICIDE	2.0 L/ha	3 leaf to just before flag leaf
FOXXY™ HERBICIDE + CLOBBER M HERBICIDE	0.8 L/ha + 2.0 L/ha	3 leaf to just before flag leaf
FOXXY HERBICIDE + 2,4-D Ester Herbicide	0.6 L/ha + 560 g a.e./ha or 0.45 L/ha + 420 g a.e./ha	4 leaf to flag leaf
FOXXY HERBICIDE + MCPA LV500 Herbicide or MCPA ESTER 600 Herbicide	0.6 L/ha + MCPA LV500 Herbicide at 1.12 L/ha or 0.6 L/ha + MCPA Ester 600 Herbicide at 0.93 L/ha.	3 leaf to just before flag leaf
Thumper® EC	1.0 L/ha	2 leaf to early flag leaf
2,4-D Amine ³ (assume 500 series)	840 mL to 1.1 L/ha	3 leaf to flag leaf
MPCA Amine ³ (assume 500 series)	840 mL to 1.1 L/ha	3 leaf to flag leaf
MCPA Ester (assume 500 series)	840 mL to 1.1 L/ha	3 leaf to flag leaf
Pardner®	1.0 L/ha	2 leaf to flag leaf
Ally®	7.5 g/ha	2 leaf to flag leaf
Attain™ HERBICIDE Tank Mix	600 mL/ha of Attain A + 1.0 L/ha of Attain B	4 leaf to flag leaf

¹ Always consult the label of the broadleaf herbicide prior to use.

² Addition of surfactants other than Score, ASSIST or CHEM SPRAY Crop Oil Concentrate is not required.

³ A reduction in control of green foxtail and wild oats may be observed when **AURORA HERBICIDE** is tank mixed with 2,4-D Amine and MCPA Amine.

Temporary crop injury may occur with tank-mixes under extreme weather conditions or when the crop is suffering from stress due to inadequate or abnormally high moisture level or extreme temperatures.

In some cases, tank mixing a pest control product with another pest control product or a fertilizer can result in biological effects that could include but are not limited to: reduced pest efficacy or increased host crop injury. The user should contact AgraCity Crop & Nutrition Ltd. at 1-844-269-3276 or www.agracity.com for information before mixing any pesticide or fertilizer that is not specifically recommended on this label.

TANK MIXES OF AURORA HERBICIDE WITH BROADLEAF WEED HERBICIDES - AERIAL APPLICATION:

CROP: SPRING WHEAT and DURUM WHEAT

For broad spectrum control of wild oats, green foxtail and broadleaf weeds, **AURORA HERBICIDE** can be tank-mixed with Buctril M. Consult the label of the tank-mix partner for a list of broadleaf weeds controlled, rates, timing, recropping restrictions, grazing interval restrictions, recommendations for specific weeds, directions for use and precautions and follow the more restrictive label. Use in a minimum of 30L of water per hectare.

¹ Always consult the label of the broadleaf partner prior to use.

BUFFER ZONES

Spot treatments using hand-held equipment DO NOT require a buffer zone.

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), and sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands).

Method of Application	Crop		Buffer zones (metres) required for the Protection of:		
			Freshwater Habitat of Depths:		Freshwater Habitat of Depths:
			Less than 1 m	Greater than 1 m	
Field sprayer	Spring and durum wheat		1	0	1
Aerial	Spring and durum wheat (55.2-70.2 g a.i./ha)	Fixed wing	1	0	20
		Rotary wing	1	0	20

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Buffer Zone Calculator in the Pesticides portion of Canada.ca.

MIXING INSTRUCTIONS - GROUND APPLICATION:

- 1 Clean spray tank and half fill with clean water. Start agitation or bypass system
- 2 If a broadleaf herbicide, insecticide or fungicide is to be used, add the product **FIRST** prior to adding **AURORA HERBICIDE** and agitate for 2-3 minutes.
- 3 Add correct amount of **AURORA HERBICIDE**.
- 4 Agitate for 2-3 minutes.
- 5 Add correct amount of either SCORE, ASSIST or CHEM SPRAY Crop Oil Concentrate.
- 6 Agitate for 1-2 minutes before adding remainder of water and then maintain constant agitation.
- 7 After any break in spraying operations, agitate thoroughly before spraying again.
- 8 Use the spray suspension as soon as it is prepared.
- 9 If an oil film starts to build up in the tank, drain tank and then clean with a detergent.

SPRAYING INSTRUCTIONS - GROUND APPLICATION:

- 1 **AURORA HERBICIDE** can be applied by ground or air. For aerial application instructions, refer to the section entitled "SPRAYING INSTRUCTIONS - AERIAL APPLICATION" which follows this section.
- 2 Water Volume: 50 to 100 litres per hectare when applied alone and a minimum of 100L/ha when tank-mixed with broadleaf herbicides
- 3 Spray Nozzles: 80° or 110° flat fan stainless steel nozzles are recommended for optimal spray coverage. Application of the spray mixture at a 45° angle in the direction of travel will result in improved spray coverage. Use 50 mesh nozzle screens. Do not use flood type nozzles, controlled

PMRA Approved Label – M.A.
2021-4039 2021-09-27

- droplet application equipment, spray foils or hollow cone nozzles.
- 4 Pressure: 275-310 kPa.
 - 5 Apply uniformly at 6-8 km/hr and avoid overlapping. Shut off spray boom while starting, turning, slowing or stopping to prevent crop injury from an over application.

Field sprayer application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Boom height must be 60 cm or less above the crop or ground.

AERIAL APPLICATION

Generic Aerial Application Label Instructions - Directions for Use

Apply only by fixed-wing or rotary aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices.

Use Precautions

Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the National Aerial Pesticide Application Manual, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.

Operator Precautions

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

During mixing, loading, clean-up, repair and when applying by groundboom, workers must wear coveralls over long-sleeved shirt and long pants, shoes and socks, and chemical-resistant gloves. When applying by aerial application, pilots must wear long-sleeved shirt and long pants.

Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the existing ground boom label.

All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

Product Specific Precautions

Read and understand the entire label before opening this product. If you have questions, call AgraCity Crop & Nutrition Ltd. at 1-844-269-3276.

Application of this specific product must meet and/or conform to the following:

MIXING INSTRUCTIONS - AERIAL APPLICATION:

1. Fill the mixing tank 1/2 full with clean water. Start gentle agitation.
2. If a broadleaf herbicide is to be used, add the product **FIRST** prior to adding **AURORA HERBICIDE** and agitate for 2-3 minutes then add correct amount of **AURORA HERBICIDE**.

PMRA Approved Label – M.A.
2021-4039 2021-09-27

3. Agitate for 2-3 minutes.
4. Add correct amount of an adjuvant of either Score, Assist or CHEM SPRAY Crop Oil Concentrate.
5. Agitate for 1-2 minutes before adding remainder of water and then maintain constant agitation.
6. Fill aircraft spray tank and maintain gentle agitation while spraying.
7. After any break in spraying operations, agitate thoroughly before spraying again. Do not let contents stand without agitation.
8. Use the spray suspension as soon as it is prepared.
9. If an oil film starts to build up in the tank, drain tank and then clean with a detergent.

SPRAYING INSTRUCTIONS - AERIAL APPLICATION:

- 1 **AURORA HERBICIDE** can be applied by ground or by air. For ground application instructions, refer to the section entitled "SPRAYING INSTRUCTIONS - GROUND APPLICATION" which precedes this section.
- 2 Water Volume - Aerial application: Minimum of 30 litres per hectare when applied alone or when tank mixed with Buctril M.
- 3 Ensure uniform application. To avoid uneven or overlapped application, use appropriate marking devices. Do not use human flaggers.

Aerial application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply when wind speed is greater than 16 km/h at flying height at the site of application. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) coarse classification. Reduce drift caused by turbulent wingtip vortices. The nozzle distribution along the spray boom length MUST NOT exceed 65% of the wing- or rotorspan.

SPRAYER CLEAN-UP:

- 1 Thoroughly clean application equipment immediately after spraying. Ensure that all traces of the product are removed. The following recommendations are provided:
- 2 Drain and flush tank walls, boom and all hoses for ten minutes with clean water. Do not clean the sprayer near desirable vegetation, wells, or other water sources.
- 3 Remove the nozzles and screens and wash separately.
- 4 Dispose of all rinsings in accordance with provincial regulations.
- 5 If a broadleaf tank-mix partner is used, always check tank-mix partner label for any additional clean up procedures.

Resistance-Management Recommendations

For resistance management, **AURORA HERBICIDE** is a Group 1 herbicide. Any weed population may contain or develop plants naturally resistant to **AURORA HERBICIDE** and other Group 1 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of **AURORA HERBICIDE** or other Group 1

PMRA Approved Label – M.A.
2021-4039 2021-09-27

- herbicides with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted.
- HERBICIDE use should be based on an IPM program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical), cultural, biological and other chemical control practices.
- Monitor treated weed populations for resistance development.
- Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment and planting clean seed.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotype.
- For further information or to report suspected resistance, contact AgraCity Crop & Nutrition Ltd at 1-844-269-3276 or at www.AgraCity.com.

NOTE TO USER:

READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR THE INDICATED SPECIAL USE APPLICATIONS:

The DIRECTIONS FOR USE for the uses described in this section of the label were developed by persons other than NewAgco, Inc. under the User Requested Minor Use Label Expansion program. For these uses, NewAgco, Inc. has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) under all environmental conditions or for all crop varieties when used in accordance with the label. The user should test the product on a small area first, under local conditions and using standard practices, to confirm the product is suitable for widespread application.

DIRECTIONS FOR USE IN THE OKANAGAN AND CRESTON FLATS REGION OF BC:

AURORA HERBICIDE can be used for the control of wild oats, green foxtail and yellow foxtail in spring wheat and durum wheat in the Okanagan and Creston Flats Regions of British Columbia. For information on crop and weed stages, rates of application, mixing and spraying instructions and precautions see the appropriate sections elsewhere on this label.

AURORA®, CANUCK®, CLOBBER®, FOXXY™, and MPOWER® are trademarks of AgraCity Crop and Nutrition Ltd. or its affiliates.

All other products mentioned are trademarks of their respective holders.