

GROUP
2,4

AgraCity

PERFORMANCE
100%
GUARANTEE

BATTLESTAR®

WHY USE BATTLESTAR?

- Same active as Stellar
- Powerful control of cleavers, and hemp-nettle, lamb's quarters, and wild buckwheat
- Controls or suppresses 24 broadleaf weeds
- Superior control of group 2 resistant cleavers
- Use in all soil zones
- 3 different actives with systemic mode of action
- Wide window of application
- Excellent crop safety
- Great tank mix with **Himalaya™**
- No re-cropping restrictions
- Superior broad-spectrum broadleaf weed control at the best price

USE ON

- Spring wheat, durum wheat, barley

WEEDS CONTROLLED

- Burdock¹
- Cleavers
- Cocklebur
- Plantain
- Lamb's-quarters
- Mustards (except dog and tansy)
- Prickly lettuce
- Ragweeds
- Stinkweed
- Vetch
- Wild radish
- Wild (annual) sunflower

¹ Before the 4-leaf stage

WEEDS SUPPRESSED

- Flixweed
- Russian pigweed
- Shepherd's-purse

SAME ACTIVE AS STELLAR™

Florasulam + Fluroxypyr + MCPA Ester 600 Herbicide

REQUIRED SURFACTANT

- None required

ACTIVE INGREDIENT

- Florasulam 50 g/L soluble concentrate (SC)
- Fluroxypyr 180 g/L emulsifiable concentrate (EC)
- MCPA Ester 600 g/L emulsifiable concentrate (EC)

PACKAGE SIZES

- **Case:** 40 acres

WHEN TO APPLY

- **Crop stage:** 3 leaf - just before flag leaf
- **Weed stage:** 1 – 4 leaf

WATER VOLUME

- 10 US gallons/acre (40 L/acre)

RAINFAST PERIOD

- 2 hours

RE-CROPPING OPTIONS

- Grow all major crops the year after application

SUPPORTED TANK MIXES*

- **Himalaya**, Axial®, Simplicity™

STORAGE

- Store in dry heated area; if product is frozen, bring to room temp and agitate before use.

GRAZING AND HARVEST

- Allow 7 days after application before cutting hay or harvesting forage.
- Mature crop may be harvested 60 days after application.

MIXING INSTRUCTIONS

1. Fill the spray tank 1/2 full of water, begin agitation, and continue throughout mixing and application process.
2. Add required rate of any dry formulation herbicide (such as **Himalaya** or Simplicity) and dissolve.
3. Add the required rate of **Battlefront** and dissolve.
4. If required, add the correct of liquid grass weed control herbicide.
5. Add the required rate of **Foxyy**™, followed by MCPA Ester.
6. Add any required surfactant, complete filling tank and apply.
7. If application is stopped prior to completion, re-suspend spray solution by full agitation prior to commencing spraying again.

UNDERSTANDING PERFORMANCE

- For optimal performance apply between 12° and 24° C in warm moist growing conditions.
- Typical symptoms (discolouration) of dying weeds may not be noticeable for 1 to 2 weeks after application, depending upon growing conditions and weed susceptibility. Control of weeds may take 3 to 4 weeks.
- Degree of control and duration of effect are dependent on weed sensitivity, weed size, crop competition, growing conditions at and following treatment, and spray coverage.
- Weeds hardened off by cold weather or drought stress may not be adequately controlled or suppressed and re-growth may occur.
- If weeds are stressed due to drought, flooding or prolonged hot or cool temperatures less than 8° C or great than 27° C, control can be reduced or delayed since weeds are not actively growing.
- Do not make applications to weeds stressed longer than 20 days due to lack of moisture as unsatisfactory control can result.

Always read and observe all label directions including rates, restrictions, and grazing and harvest limitations.

*Supported by AgraCity under PMRA tank mix guidelines. When applying tank mix combinations, read and observe all label directions for each product used in the tank mix, and follow the more stringent label precautionary measures for mixing, loading and applying stated on both product labels.