

TAIPAN[®] HERBICIDE

AGRACITY EXCLUSIVE PRODUCT Bentazon + Imazamox Herbicide

WHY USE TAIPAN?

- AgraCity exclusive product - excellent control of many broadleaf weeds
- Proven control of cocklebur, cow cockle, green smartweed, sheperd's purse, stinkweed, volunteer canola and grassy weeds
- Controls or suppresses 23 broadleaf and grass weeds
- Use in all soil zones
- Multiple modes of action for resistance management
- Early weed control timing for higher yields
- Fewer re-cropping restrictions than some Group 2 only herbicide products
- Tank mix options for improved grass control
- Trusted broad-spectrum control of grass and broadleaf weeds

FOR USE ON

- Field peas



WEEDS

CONTROLLED

Grass Weeds

- 1-4 main leaf stage:
- Barnyard grass
 - Green foxtail
 - Volunteer barley
 - Volunteer wheat¹
 - Volunteer tame oats
 - Wild oats

Broadleaf Weeds

- Cotyledon-4 leaf stage
- Bird rape
 - Cocklebur
 - Cow cockle
 - Flower of an hour
 - Green smartweed
 - Lady's thumb
 - Lamb's-quarters
 - Redroot pigweed
 - Russian thistle²
 - Shepherd's purse
 - Stinkweed
 - Velvetleaf
 - Volunteer canola
 - Wild mustard

SUPPRESSED

- Cleavers
- Japanese brome
- Kochia²
- Round-leaved mallow
- Wild buckwheat

¹ Not including Clearfield® wheat

² Not including Group 2 resistant biotypes



REQUIRED SURFACTANT

- 28% UAN at 0.81 L/ac plus Merge® at 0.5% v/v

ACTIVE INGREDIENTS

- Imazamox 70% water dispersible granule (WDG)
- Bentazon 480 g/L solution

PACKAGE SIZES

- **Case:** 40 acres
- **Bulk:** 960 acres

WHEN TO APPLY

- Peas: 3-6 above ground nodes

WATER VOLUME

- Minimum 10 US gallons/acre (40 L/acre)

RAINFAST PERIOD

- 6 hours

RE-CROPPING OPTIONS

- **3 months after application:** Winter wheat
- **1 year after application:** barley, canary seed, canola*, chickpeas, corn, durum wheat, peas, flax, lentils, soybeans, spring wheat, sunflowers, tame oats.
- **2 years after application:** the above crops plus mustard (condiment type only)
- A field bioassay is recommended before growing crops other than those listed above.

TANK MIX OPTIONS*

- Quiz®

STORAGE

- Do not freeze.

GRAZING AND HARVEST

- Do not graze treated pea fields or cut for feed within 20 days of application.
- Do not apply within 60 days of harvest.

MIXING INSTRUCTIONS

1. Fill spray tank 3/4 full of water, start sprayer agitation and continue throughout mixing and spraying process.
2. Add the required amount of **Samurai®** and agitate until dissolved.
3. Add the required amount of **Boa®**.
4. Add the required amount of grass tank mix partner if desired.
5. Add required amount of 28% UAN.
6. Add the required amount of Merge surfactant.
7. Add the remaining amount of water and apply.
8. If application is interrupted prior to completing application, re-suspend spray solution by agitating prior to resuming application
9. Check in-line and nozzle screens between loads and clean if necessary.

UNDERSTANDING PERFORMANCE

- Optimum control occurs when applied between 12°C and 24° C in warm moist growing conditions.
- Thorough coverage is critical — use a minimum water volume of 10 gallons/acre.
- Symptoms of control (discoloration at the growing point) will take several hours and several days to be visible, depending on growing conditions. Control of weeds may take 3-4 weeks.
- Do not spray **Taipan** if temperatures of 5° C or less are forecast within 3 days of application. Application just before or after cold temperatures can result in crop injury and reduced weed control.
- Drought, crop stress, and cold or hot temperatures will negatively affect performance as weeds are not actively growing.
- Do not apply **Taipan** to crops that have been subjected to stress from conditions such as hail damage, flooding, drought, hot, humid weather, widely fluctuating temperature conditions, prolonged cold conditions or injury from prior herbicide applications as this may lead to poor results and/or crop damage.
- Occasionally transient crop yellowing may be observed after application, but this is temporary and will be outgrown and should not affect yield.

Always read and follow all label precautions and directions.

*Initial injury to non-Clearfield® canola may be observed

*When applying tank mix combinations, always follow all label requirements for each product used in the tank mix. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed.

Boa®, Quiz®, and Samurai® are trademarks of AgraCity or its affiliates. All other products are trademarks of their respective holders.